

Susceptible–infected–recovered epidemics in dynamic contact networks

Erik Volz* and Lauren Ancel Meyers

Department of Integrative Biology, University of Texas at Austin, 1 University Station, C0930, Austin, TX 78712, USA

Contact patterns in populations fundamentally influence the spread of infectious diseases. Current mathematical methods for epidemiological forecasting on networks largely assume that contacts between individuals are fixed, at least for the duration of an outbreak. In reality, contact patterns may be quite fluid, with individuals frequently making and breaking social or sexual relationships. Here, we develop a mathematical approach to predicting disease transmission on dynamic networks in which each individual has a characteristic behaviour (typical contact number), but the identities of their contacts change in time. We show that dynamic contact patterns shape epidemiological dynamics in ways that cannot be adequately captured in static network models or mass-action models. Our new model interpolates smoothly between static network models and mass-action models using a mixing parameter, thereby providing a bridge between disparate classes of epidemiological models. Using epidemiological and sexual contact data from an Atlanta high school, we demonstrate the application of this method for forecasting and controlling sexually transmitted disease outbreaks.

Keywords: infectious disease; susceptible–infected–recovered; networks; concurrency; syphilis

1. INTRODUCTION

Most epidemic models incorporate a homogeneous mixing assumption, sometimes called the law of mass action (Ross 1910; Anderson & May 1991; Diekmann & Heesterbeek 2000), whereby the rate of increase in epidemic incidence is proportional to the product of the number of infectious and susceptible individuals. This assumption has been relaxed in some compartmental (van den Driessche & Watmough 2002) and metapopulation models (Lloyd & May 1996; Finkenstadt & Grenfell 1998; Grenfell *et al.* 2001; Watts *et al.* 2005), but not eliminated. The mass-action assumption is robust in the sense that it is consistent with several scenarios for the individual-to-individual transmission of disease. In particular, it is equivalent to a model in which all individuals in a population make contact at an identical rate and have identical probabilities of disease transmission to those contacts per unit of time. Although this assumption is unrealistic, it facilitates mathematical analysis and, in some cases, offers a reasonable approximation.

Populations can be quite heterogeneous with respect to susceptibility, infectiousness, contact rates or number of partners, and simple homogeneous mixing models do not allow for extreme variation in host parameters. New network-based mathematical methods capture some, but not all, aspects of population heterogeneity (Gupta *et al.* 1989; Andersson 1998; Callaway *et al.* 2000; Liljeros *et al.* 2001; Strogatz 2001; Newman 2002a; Newman *et al.* 2002). Ideally, an epidemic model would incorporate the following realities of human-to-human contacts:

— A given individual has contact with only a finite number of other individuals in the population at any one time, and

contacts that can result in disease transmission are usually short and repeated events.

- The number and frequency of contacts between individuals can be very heterogeneous.
- The numbers and identities of an individual's contacts will change as time goes by.

The first two points have been addressed previously by static network models (Eames & Keeling 2002; Newman 2002a; Meyers *et al.* 2005, 2006). We will focus on the third point and introduce a modelling framework that allows an individual's contacts to change in time.

Concurrent and serial contacts were first shown to be important to HIV transmission dynamics by Dietz (1988) and Watts & May (1992). In the public health literature, the number of concurrent contacts has long been understood to be important for sexually transmitted disease (STD) transmission (Ford *et al.* 2002; Manhart *et al.* 2002; Adimora *et al.* 2006), independently of cumulative number of contacts. Epidemics in dynamic networks have been modelled using high-dimensional pair-approximation methods (Altmann 1995, 1998; Eames & Keeling 2004) and moment closure methods on dynamic contact networks (Ferguson & Garnett 2000; Bauch 2002). Others have conducted simulation-based studies (Chick *et al.* 2000; Doherty *et al.* 2006).

Here, we introduce a low-dimensional system of non-linear ordinary differential equations (ODEs) to model susceptible–infected–recovered (SIR) epidemics in a simple class of dynamic network. We find this provides a useful departure point for mathematical analysis, although it does not incorporate as much detail as some other models. This model complements earlier work by providing a relatively simple framework that is less computationally demanding, easier to implement and amenable to the derivation of fundamental results. Our methods are compared with the

* Author for correspondence (erik.volz@mail.utexas.edu).

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2007.1159> or via <http://www.journals.royalsoc.ac.uk>.

more commonly used pair-approximation methods in the electronic supplementary material.


We use our dynamic network model to characterize the impacts of population heterogeneity and contact rates on epidemic dynamics. We further show that the model reproduces basic classes of epidemiological models, such as the standard mass-action SIR model and static network model, as the parameter that controls population mixing varies.

2. NEIGHBOUR EXCHANGE MODEL

Human contact patterns are potentially complex, as the numbers and intensity of contacts can vary considerably across a population. Furthermore, contacts are transitory events such that the identities of one's contacts change in time. To capture such heterogeneity, we introduce the *neighbour exchange* (NE) model as a simple extension of a static contact network model. In this model, an individual's number of concurrent contacts remains fixed while the composition of those contacts changes at a specified rate. The model assumes that at any given time, an individual will be in contact with an individual-specific number of neighbours with whom disease transmission is possible. Each contact is temporary, lasting a variable amount of time before coming to end, at which point the neighbour is replaced by a different individual.

Let the population of interest consist of n individuals, each of which falls into one of three exclusive states: susceptible, infectious or recovered. At some time t , an individual ego will have k_{ego} contacts with other individuals (i.e. *alters*): $(ego, alter_1), (ego, alter_2), \dots, (ego, alter_k)$. Only undirected contact networks will be considered such that if there exists a contact $(ego, alter)$ there will also be a contact $(alter, ego)$. In network terminology, a directed link, denoted $(ego, alter)$, is called an *arc*. An undirected link, denoted $\{ego, alter\}$, is called an *edge*. The *degree* of a node ego is the number of edges connected to the node. The term *contact* will be used specifically to denote a directed arc in the network, where two arcs correspond to each undirected edge. The *k-degree* of a node will be the number of concurrent contacts to/from the node (table 1).

The NE model assumes that the identities of a node's neighbours will continually change, while the total number of current neighbours remains constant. This occurs through an exchange mechanism in which the destination nodes of two edges are swapped. For example, two nodes ego and ego' with distinct contacts $(ego, alter)$ and $(ego', alter')$ may exchange contacts such that these are replaced with $(ego, alter')$ and $(ego', alter)$. There are two edges and four contacts involved in each edge swap. The fate of each edge and contact is summarized in the following pseudo-chemical equation:



In the model, any given contact $(ego, alter)$ will be reassigned to $(ego, alter')$ at a constant rate ρ . Equivalently, a given edge is swapped at a rate $\rho/2$.

A node's degree never changes during an edge swap and thus the degree distribution is preserved. This property has

Table 1. Notation for epidemic and network parameters.

S	the fraction of population susceptible
I	the fraction of population infectious
R	the fraction of population recovered
\mathcal{S}	set of susceptible nodes
\mathcal{I}	set of infectious nodes
\mathcal{R}	set of recovered nodes
\mathcal{J}	cumulative epidemic incidence ($\mathcal{J} = I + R$)
r	transmission rate
μ	recovery rate
ρ	mixing rate
k_{ego}	degree of node ego
p_k	the fraction of nodes with degree k

Table 2. Notation for network and epidemic quantities.

\mathcal{A}_X	set of contacts $(ego, alter)$ s.t. $ego \in X$
\mathcal{A}_{XY}	set of contacts $(ego, alter)$ s.t. $ego \in X$ and $alter \in Y$
M_X	fraction of contacts in set \mathcal{A}_X
M_{XY}	fraction of contacts in set \mathcal{A}_{XY}
$p_I = M_{SI}/M_S$	fraction of contacts from susceptibles which go to infectious nodes
$p_S = M_{SS}/M_S$	fraction of contacts from susceptibles which go to other susceptible nodes

made the random edge swap a common starting point for investigations of dynamic networks. Discrete-time versions of the rewiring process have been well studied in applications outside of epidemiology (Watts & Strogatz 1998; Maslov & Sneppen 2002; Evans & Plato 2007; Holme & Zhao 2007)

For mathematical tractability, we make a few simplifying assumptions about the epidemic process. First, for the duration of a contact, infectious individuals transmit disease to neighbours at a constant rate r . Second, infectious individuals become recovered at a constant rate μ .

We also simplify the mathematics by considering only the simplest category of heterogeneous networks—semi-random networks that are random with respect to a given degree distribution (Molloy & Reed 1995, 1998). The degree distribution will have density p_k , which is the probability that a node chosen uniformly at random has k concurrent contacts.

The simultaneous epidemic and network dynamics described above collectively determine the NE model. An even more realistic model would allow the number k of concurrent contacts of a node to vary stochastically, but the current model offers a valuable first step towards understanding epidemiological processes on dynamic host networks.

(a) Dynamics

We will expand on the dynamic probability generating function (PGF) techniques first introduced in Volz (2007) to model SIR-type epidemics in static networks. These techniques are powerful and are easily extended to consider dynamic contact networks. We start with an overview of the basic model and then introduce additional terms that model the NE process.

The concurrent degree distribution p_k will be generated by the PGF

$$g(x) = p_0 + p_1x + p_2x^2 + p_3x^3 + \dots \tag{2.2}$$

The dummy variable x in this equation serves as a placeholder for dynamic variables.

Let \mathcal{A}_X be the sets of contacts (arcs) where *ego* is in the set X . We will consider the sets of all susceptible, infected and recovered nodes, denoted $X = S$, $X = I$ and $X = R$, respectively. $M_X = \#\{\mathcal{A}_X\}/\#\{A\}$ is then defined as the fraction of total contacts in set \mathcal{A}_X . Now let \mathcal{A}_{XY} be the set of contacts such that *ego* is in the set X and *alter* is in the set Y , and $M_{XY} = \#\{\mathcal{A}_{XY}\}/\#\{A\}$ be the fraction of total contacts in set \mathcal{A}_{XY} . For example, M_{SS} is the fraction of all arcs in the network that connect two currently susceptible nodes (table 2).

The following derivation assumes that each contact of a susceptible node (*ego*, *alter*) $\in \mathcal{A}_S$ has a uniform probability that *alter* $\in \mathcal{I}$, denoted $p_I = M_{SI}/M_S$, and a uniform probability $p_S = M_{SS}/M_S$ that *alter* $\in S$. A degree k susceptible node has an expected number kp_I contacts with infectious nodes; and, in a small time dt , an expected number $rkp_I dt$ of a degree k susceptible nodes' contacts will transmit disease to that node. The instantaneous hazard of infection for a degree k susceptible node is then given by

$$\lambda_k(t) = rkp_I(t). \tag{2.3}$$

Let $u_k(t)$ denote the fraction of degree k nodes remaining susceptible at time t . Equation (2.3) implies

$$\begin{aligned} u_k(t) &= \exp\left\{\int_{\tau=0}^t -\lambda_k(\tau)d\tau\right\} \\ &= \exp\left\{\int_{\tau=0}^t -rkp_I(\tau)d\tau\right\} \\ &= \exp\left\{\int_{\tau=0}^t -rp_I(\tau)d\tau\right\}^k. \end{aligned} \tag{2.4}$$

Now let $\theta = u_1(t)$ be the fraction of degree $k=1$ nodes in the network that remain susceptible at time t . Equation (2.4) implies that $u_k = \theta^k$. (Henceforth, variables that are clearly dynamic, for example θ , appear without a time (t) variable.)

We use the PGF for the network degree distribution (equation (2.2)) to calculate the fraction of nodes that remain susceptible at time t .

$$S = p_0 + u_1p_1 + u_2p_2 + \dots = \sum_k p_k\theta^k = g(\theta). \tag{2.5}$$

The dynamics of θ in equation (2.5) are given by

$$\dot{\theta} = \frac{d}{dt}u_1 = -\lambda_1\theta = -rp_I\theta. \tag{2.6}$$

Unfortunately, this does not form a closed system of differential equations, as equation (2.6) depends on the dynamic variable p_I . From the definition of p_I we have

$$\dot{p}_I = \frac{d}{dt}\frac{M_{SI}}{M_S} = \frac{\dot{M}_{SI}}{M_S} - \frac{M_{SI}\dot{M}_S}{M_S^2}. \tag{2.7}$$

To obtain the derivatives of M_{SI} and M_S , we observe that, in time dt , $-\dot{S}$ nodes become infectious, resulting in modifications to the sets \mathcal{A}_{SI} , \mathcal{A}_{IS} and \mathcal{A}_S . In particular, any given arc from a newly infected individual was formerly in one of \mathcal{A}_{SS} , \mathcal{A}_{SI} or \mathcal{A}_{SR} and is now in one of \mathcal{A}_{IS} , \mathcal{A}_{II} or \mathcal{A}_{IR} . The $-\dot{S}$ new infected are not selected uniformly at random from the susceptible population, but rather with

probability proportional to the number of contacts to infectious nodes.

We pause for a few definitions. First, it is useful to break the degree of a node into three quantities: the number of contacts to currently susceptible, infected and recovered nodes. We refer to these as X -degrees, where $X = S, I$ or R , respectively. Second, imagine following a randomly chosen contact to its *alter* and counting all the edges emanating from that node, except for the one on which we have arrived. We call the resulting total quantity the *excess degree* of the node, and the resulting neighbour-specific quantities *excess X-degrees* of the node, where X indicates one of the three possible disease states.

We introduce the notation $\delta_{X,Y}(Z)$ to represent the average excess Z -degree of nodes currently in disease state X selected by following a randomly chosen arc from the set \mathcal{A}_{YX} ; in other words, the excess Z -degree of a node of type X selected with probability proportional to the number of contacts to nodes of type Y . We further define $\delta_{X,Y}$ to be the average (total) excess degree of nodes currently in disease state X selected by following a randomly chosen arc from the set \mathcal{A}_{YX} . For example, imagine first randomly choosing an arc from \mathcal{A}_{IS} , then following that arc to its destination (susceptible) node and finally counting all of the other edges emanating from that node (ignoring the one along which we arrived). Then $\delta_{S,I}(S)$, $\delta_{S,I}(I)$ and $\delta_{S,I}(R)$ give the average number of contacts to other susceptible, infected and recovered nodes chosen in this way, respectively; $\delta_{S,I}$ gives the average total number of contacts emanating from nodes chosen in this way.

Using this notation, the equations for \dot{M}_{SI} and \dot{M}_S are as follows (for more details, see Volz (2007))

$$\dot{S} = \frac{d}{dt}g(\theta) = \dot{\theta}g'(\theta) = -rp_I\theta g'(\theta), \tag{2.8}$$

$$\dot{M}_{SI} = ((-\dot{S})\delta_{S,I}(S) - (-\dot{S})\delta_{S,I}(I))/g'(1) - (r + \mu)M_{SI}, \tag{2.9}$$

$$\dot{M}_{SS} = -2(-\dot{S})\delta_{S,I}(S)/g'(1) \quad \text{and} \tag{2.10}$$

$$\dot{M}_S = \frac{d}{dt}\theta g'(\theta)/g'(1) = -(rp_I\theta g'(\theta) + rp_I\theta^2 g''(\theta))/g'(1). \tag{2.11}$$

The calculations of the $\delta_{X,Y}(Z)$ are straightforward and based on the current degree distribution of susceptible nodes. The calculations are given in the electronic supplementary material and in Volz (2007) and result in the following:

$$\delta_{S,I}(I) = p_I\theta g''(\theta)/g'(\theta) \quad \text{and} \tag{2.12}$$

$$\delta_{S,I}(S) = p_S\theta g''(\theta)/g'(\theta). \tag{2.13}$$

Combining the equations for M_S , \dot{M}_S , and \dot{M}_{SI} yields the dynamics of p_I in terms of the parameters r and μ , the PGF $g(\cdot)$, and the dynamic variable p_S . The resulting model is given in table 3. The dynamics for p_S complete the model and can be derived analogously to the equation for p_I (see the electronic supplementary material for details).

(i) *Dynamic contact networks*

We now extend the model given in table 3 to allow NEs at a rate ρ . First consider θ . An edge swap (equation (2.1)) will affect the arrangement of edges among susceptible,

Table 3. System of equations used to model the spread of SIR epidemic in a static semi-random network.

$$\begin{aligned} \dot{\theta} &= -r p_I \theta \\ \dot{p}_S &= r p_S p_I (1 - \theta g(\theta)/g'(\theta)) \\ \dot{p}_I &= r p_I p_S \theta g(\theta)/g'(\theta) - p_I (1 - p_I) r - p_I \mu \\ S &= g(\theta) \end{aligned}$$

infectious or recovered nodes; however, it will never directly cause a node to change its disease state. The dynamics of θ are only indirectly influenced by NE dynamics, through changes to p_I , and thus equation (2.6) still holds.

NE does, however, affect the composition of contacts between susceptible and infectious nodes. We postulate that the equations for \dot{p}_I and \dot{p}_S can be expressed in the following modified forms:

$$\dot{p}_I = r p_I p_S \theta g''(\theta)/g'(\theta) - p_I (1 - p_I) r - p_I \mu + f_I(p_I, M_I) \tag{2.14}$$

and

$$\dot{p}_S = r p_S p_I \left(1 - \theta \frac{g''(\theta)}{g'(\theta)} \right) + f_S(p_S, M_S), \tag{2.15}$$

where the functions $f_x(\cdot, \cdot)$ represent the contribution of NE dynamics to the system.

NE dynamics can both decrease or increase the values of p_I and p_S . First, consider the decrease of p_I due to NE dynamics.

- At rate ρ , a given contact (*ego*, *alter*) will transform to (*ego*, *alter'*).
- Given that (*ego*, *alter*) $\in \mathcal{A}_S$, *alter* $\in \mathcal{I}$ with probability p_I .
- With probability $1 - M_I$, *alter'* $\notin \mathcal{I}$.

Thus p_I will be decreased by NE dynamics at a rate $\rho p_I (1 - M_I)$.

Similarly, p_I can increase due to NE as follows:

- At rate ρ , a given contact (*ego*, *alter*) will transform to (*ego*, *alter'*).
- Given that (*ego*, *alter*) $\in \mathcal{A}_S$, *alter* $\notin \mathcal{I}$ with probability $1 - p_I$.
- With probability M_I , *alter'* $\in \mathcal{I}$.

Thus p_I is increased by NE dynamics at rate $\rho (1 - p_I) M_I$. We add the positive and negative contributions to calculate the total influence of NE on p_I .

$$f_I(p_I, M_I) = \rho (M_I - p_I). \tag{2.16}$$

By similar reasoning, we determine $f_S(p_S, M_S)$.

$$\begin{aligned} f_S(p_S, M_S) &= \rho (\Pr[\textit{alter} \notin \mathcal{S}] \times \Pr[\textit{alter}' \in \mathcal{S}] \\ &\quad - \Pr[\textit{alter} \in \mathcal{S}] \times \Pr[\textit{alter}' \notin \mathcal{S}]) \\ &= \rho (M_S - p_S) = \rho ((\theta)g'(\theta)/g'(1) - p_S). \end{aligned} \tag{2.17}$$

The complete system of NE-adjusted equations is reported in table 4. The dynamic variable M_I , however, appears in equation (2.17) and cannot be put in terms of the variables θ , p_S and p_I . Therefore, a fourth dynamical equation must be included for M_I , which is listed in table 4. Fortunately, the dynamics are very straightforward. Recall that $\delta_{S,I}$ denotes the average excess degree of susceptibles chosen by following randomly selected arcs between infectious and susceptible nodes, and thus indicates the average excess degree for individuals who become infected in a small time interval dt . We add one ($\delta_{S,I} + 1$) to obtain

Table 4. Deterministic NE model. (This system of equations is used to model the spread of an SIR-type epidemic in a dynamic semi-random network with stochastic exchange of neighbours at constant rate ρ .)

$$\begin{aligned} \dot{\theta} &= -r p_I \theta \\ \dot{p}_S &= r p_S p_I (1 - \theta g''(\theta)/g'(\theta)) + \rho ((\theta)g'(\theta)/g'(1) - p_S) \\ \dot{p}_I &= r p_I p_S \theta g''(\theta)/g'(\theta) - p_I (1 - p_I) r - p_I \mu + \rho (M_I - p_I) \\ \dot{M}_I &= -\mu M_I + r p_I (\theta^2 g''(\theta) + \theta g'(\theta))/g'(1) \\ S &= g(\theta) \end{aligned}$$

the average degree for such newly infected nodes. Then, recalling that M_I decays at rate μ , we have

$$\dot{M}_I = (-\dot{S})(\delta_{S,I} + 1)/g'(1) - \mu M_I. \tag{2.18}$$

All of the results that follow will assume that a very small fraction ϵ of nodes is initially infected, and thus there is only a very small probability that two initially infected individuals contact each other. Then we anticipate the following initial conditions (Volz 2007):

$$\theta = M_I = 1 - \epsilon, \tag{2.19}$$

$$p_I = \epsilon/(1 - \epsilon) \quad \text{and} \tag{2.20}$$

$$p_S = (1 - 2\epsilon)/(1 - \epsilon). \tag{2.21}$$

(b) Convergence to a mass-action model

In the limit of large mixing rate ($\rho \rightarrow \infty$), the probability of being connected to a susceptible, infectious or recovered node is directly proportional to the number of edges emanating from nodes in each state. Referring to table 4, it is clear that p_I converges instantly to M_I and p_S converges instantly to M_S . Here, we show that as the mixing rate grows, the underlying network structure becomes irrelevant and the model converges to a mass-action model.

To see this, we replace every occurrence of the variable p_I in the system of equations in table 4 with M_I . Then

$$\dot{\theta} = -r M_I \theta \quad \text{and} \tag{2.22}$$

$$\dot{M}_I = \frac{r M_I}{g'(1)} (\theta g'(\theta) + \theta^2 g''(\theta)) - \mu M_I. \tag{2.23}$$

Neither equations (2.22) or (2.23) depend on p_S , and thus together form a closed system of equations that describe the epidemic dynamics. These equations incorporate arbitrary heterogeneity in contact rates, but no longer consider an explicit contact network. When we assume that contact rates are homogeneous throughout the population, then these equations are equivalent to a simple SIR compartmental model. To illustrate, we retrieve the standard SIR dynamics by setting $g(x) = x$, which means that every node has exactly one concurrent contact. In such a population, the number of arcs to infectious individuals is exactly equal to the number of infectious nodes, that is, $p_I = I$. Then, substituting into equations (2.22) and (2.23), we reproduce the standard equations

$$S = g(\theta) = \theta, \tag{2.24}$$

$$\dot{\theta} = -r I \theta = -r I S \quad \text{and} \tag{2.25}$$

$$\dot{M}_I = r M_I \theta - \mu M_I = r I S - \mu I. \tag{2.26}$$

Equations (2.22) and (2.23) are potentially extremely useful, as they incorporate arbitrary heterogeneity into a system of equations no more complex than the standard compartmental SIR model.




Figure 1. Trajectories of the NE model (solid lines, table 4) are compared for several values of the mixing rate (ρ) with an analogous mass-action model (circles, equations (2.22) and (2.23)). The degree distribution is Poisson ($z=1.5$) and $r=\mu=0.2$.

Figure 1 demonstrates the convergence of the NE model to the corresponding mass-action model for a Poisson degree distribution ($r=\mu=0.2$, $z=1.5$). The circles indicate the solution to the mass-action model (equations (2.22) and (2.23)). We observe that the convergence is quite rapid as ρ is increased in the multiples of μ . This supports the common assumption that the mass-action model is a reasonable approximation for populations marked by many short-duration contacts.

3. STOCHASTIC SIMULATIONS

To test the NE model, we compare its predictions to stochastic simulations of an analogous epidemic process in networks. We first generate semi-random networks using the configuration model (Molloy & Reed 1995). The epidemic simulations then proceed as follows:

- (i) One node is selected uniformly at random from the population to be *patient zero*, the first infected individual.
- (ii) Each contact $a \in \mathcal{A}$ has an exchange time $\Delta t_E(a)$ drawn from an exponential distribution (parameter ρ). This time is added to a queue.
- (iii) When a node v is infected at time t , a time of infection $\Delta t_I(v, a)$ is drawn from an exponential distribution (parameter r) for each contact $a = (ego, alter)$ such that v is *ego*. The time $t + \Delta t_I(v, a)$ is added to a queue.
- (iv) When a node v is infected at time t , a time of recovery $\Delta t_R(v)$ is drawn from an exponential distribution (parameter μ) and is assigned to v . The time $t + \Delta t_R(v)$ is added to a queue.
- (v) When $t' = t + \Delta t_E(a)$ is the earliest time in the queue, an edge swap is performed, as per equation (2.1). The first edge involved in the swap corresponds to the contact a . The second edge is selected by choosing a unique element out of all edges uniformly at random. Then a new time $t' + \Delta t_E(a)$ is drawn and added to the queue.
- (vi) When $t' = t + \Delta t_I(v, a)$ is the earliest time in the queue, a transmission event will occur, providing v has not recovered. Node v transmits to whatever

node is occupying the position of *alter* at that time, causing *alter* to change state to \mathcal{I} if currently susceptible. If a transmission event occurs, a new time $t' + \Delta t_I(v, a)$ is drawn and added to the queue.

- (vii) When $t + \Delta t_R(v)$ is the earliest time in the queue, the corresponding node v enters a recovered state such that any transmission event with $v = ego$ is removed from the queue.

This process continues until there are no more transmission events in the queue.

Figures 2 and 3 show a comparison of 1000 stochastic simulations to the solution of the NE-SIR equations for two concurrent degree distributions:

- *Poisson*. $p_k = z^k e^{-z}/k!$, $k \geq 0$. This is generated by $g(x) = e^{z(x-1)}$.
- *Power law with cut-off*. $p_k = k^{-\alpha} / \sum_{i=1}^{\kappa} i^{-\alpha}$, $k \geq 1, k \leq \kappa$. This is generated by $g(x) = \sum_k k^{-\alpha} x^k / \sum_{i=1}^{\kappa} i^{-\alpha}$.

Figure 2 depicts epidemics on a network with a Poisson degree distribution ($z=1.5$) and parameters $r=0.2, \mu=0.1$ and $\rho=0.25$. Figure 3 shows epidemics on a network with a power-law degree distribution ($\alpha=2.1, \kappa=75$) and parameters $r=0.2, \mu=0.1$ and $\rho=0.20$. The deterministic NE model (table 4) predicts a trajectory that passes through the central-most region of the swarm of simulation trajectories and shows good agreement with the final size. There is nevertheless a great deal of variability among the simulation trajectories in terms of the onset of the expansion phase—the point in time when the epidemic increases at its maximal rate. At the onset of expansion phase, all trajectories are more or less similar, in agreement with the NE model.

In contrast to the homogeneous Poisson network, the power law gives an almost immediate expansion phase. This can be understood by noting that the hazard of infection is proportional to p_I , and initially

$$\dot{p}_I(t=0) = \varepsilon \left(r \frac{g''(1-\varepsilon)}{g'(1-\varepsilon)} - r - \mu - \frac{\rho\varepsilon}{1-\varepsilon} \right). \tag{3.1}$$

There is a ratio of PGFs in equation (3.1)

$$g''(1-\varepsilon)/g'(1-\varepsilon) \approx g''(1)/g'(1) = \frac{\sum_k k^2 p_k}{\sum_k k p_k} - 1.$$




Figure 2. One-thousand stochastic simulations (small dots) are compared with the predicted trajectory of a NE epidemic (large dots) in a Poisson network ($z=1.5$). $r=0.2$, $\mu=0.1$ and $\rho=0.25$.




Figure 3. One-thousand stochastic simulations (small dots) are compared with the predicted trajectory of a NE epidemic (large dots) in a power-law network ($\alpha=2.1$, $\kappa=75$). $r=0.2$, $\mu=0.1$ and $\rho=0.20$.

This is approximately the ratio of the second moment to the first moment of the degree distribution, which for the power-law approaches infinity as the cut-off $\kappa \rightarrow \infty$. Because the ratio is very large, power-law networks have almost immediate expansion phase (Pastor-Satorras & Vespignani 2001a,b; Boguna *et al.* 2003; Barthélemy *et al.* 2005).

4. APPLICATION OF THE NE MODEL TO SYPHILIS OUTBREAK AMONG ATLANTA ADOLESCENTS

We next implement the NE model using link-tracing data from a 1996 outbreak of syphilis among Atlanta adolescents. Sexual network data typically report contact time, duration of a contact and frequency of interaction. Most network models do not take into account the serial aspect of

sexual contacts and instead assume that all contacts reported in a survey are constant over the duration of an epidemic or infectious period. Here, we illustrate that the dynamic SIR network model (table 4) can explicitly capture the transitory nature of sexual contact patterns. Owing to data limitations and uncertainty around syphilis infection rates, the NE model cannot describe this particular epidemic with much fidelity. Instead, this section is simply intended to demonstrate the straightforward application of the NE model to public health data.

We use public health data from an outbreak of syphilis within an adolescent community centred on an Atlanta high school (Rothenberg *et al.* 1998). Initially, several adolescents diagnosed with syphilis were interviewed by epidemiologists. The sexual contacts of these respondents




Figure 4. The final epidemic size as predicted by the NE model (table 4) is shown with respect to the transmission rate r and recovery rate μ for the Atlanta syphilis data. Contours are also provided for $\tau = 0.637$ and 0.20 .

were then traced and interviewed. In all, 34 people were interviewed and 204 contacts were traced. Each interviewee named his/her sexual contacts and listed the date of first and last interaction with each contact.

The complexity of syphilis transmission dynamics (Garnett *et al.* 1997) and the small size of our dataset make modelling the 1996 outbreak difficult. Thus, our analysis suffers from significant uncertainty in the estimated rates and parameters, particularly the transmission and recovery rates. We offer an additional caveat that recovery from syphilis does not provide lifelong immunity; yet our model does not allow recovered individuals to return to the susceptible state. We believe that this is appropriate in so far as the transition from the infected state back to the susceptible state occurs at a low rate relative to the spread of this particular outbreak. In practice, immunity is very long due to treatment and behavioural change, such that fast-spreading epidemics can be considered in isolation without including recovery or birth into the population (Grassly *et al.* 2005).

We estimate the relevant parameters using equations given in the electronic supplementary material and in Heckathorn (2002), Salganik & Heckathorn (2004) and Volz & Heckathorn (in press). In brief, a typical syphilis infection can last about a year if left untreated, and a typical infectious period will last 154 days on average (Jones 2005). A convenient estimate of the recovery rate is then $\hat{\mu} = 1/154 = 0.0065$. This estimate ignores many features of the pathology of syphilis for mathematical convenience, such as different probabilities of recovery at different stages of the infectious period (Garnett *et al.* 1997). There are diverse estimates for the transmissibility of syphilis, ranging from 9.2 to 63% per partner (Garnett *et al.* 1997). The estimate of 63% was selected as the most credible by the authors in Garnett *et al.* (1997).

Using the contact durations reported in the Atlanta study, we estimate the mixing rate of the population to be $\hat{\rho} = 0.032$ (electronic supplementary material). We then use

the reported numbers of contacts to estimate \hat{k}_{ego} , the average number of concurrent contacts for each individual in the sample. The degree distribution p_k can then be estimated (electronic supplementary material, Heckathorn 2002; Salganik & Heckathorn 2004; Volz & Heckathorn in press) from the sequence \hat{k}_i , which is well fit by a power law with exponent $\alpha = -2.66$. For the power-law fit, $\chi^2/n = 0.018$. An exponential distribution provided a worse fit to the data with $\chi^2/n = 0.632$. We assume the estimated degree distribution in the following analysis.

Figure 4 shows the final size of an outbreak as predicted by the NE model over a broad range transmission and recovery rates. The straight contours correspond to the ratio of r to μ that yields a transmissibility $\tau = 0.637$ and 0.20 . Unfortunately, altering the transmissibility within this plausible range can yield any final size from almost 0 to 100%. From our sample, we can estimate the epidemic prevalence as 35%. We conclude that the NE model is consistent with the observed outbreak; however, this does not constitute a critical test of the suitability of the NE model to this outbreak.

5. DISCUSSION

Human contact patterns are characterized by heterogeneous number of transitory contacts. If contacts change at a rate that is slow relative to the rate of epidemic propagation, then static network approximations such as those based on bond percolation (Callaway *et al.* 2000; Newman 2002a; Meyers *et al.* 2005, 2006) may be appropriate. Alternatively, if contacts have very short duration relative to epidemic dynamics, then static network approximations break down and a mass-action model is more appropriate (equations (2.22) and (2.23)). In between these extremes, contacts are neither fixed nor instantaneous, and accurate epidemiological forecasting requires models that explicitly capture their dynamics, such as the NE model developed here. In fact, by changing a

single parameter (the mixing rate), the NE model crosses the spectrum of models from static network to mass action.

The NE model is particularly useful for building models from link-tracing data. For many datasets, standard mass-action models do not adequately capture the finite number and extended duration of contacts, whereas static network models ignore the transitory and serial nature of contacts. Using the example of a 1996 syphilis outbreak in an adolescent population, we showed that the NE model can be easily fit to sexual contact data and then used to explore the epidemiological implications of host population structure.

In the limit of large mixing rate, the NE model becomes a simple (low-dimensional) mass-action model (equations (2.22) and (2.23)) that captures SIR dynamics in populations with arbitrary heterogeneity of contact rates. It reduces to the standard mass-action model when one assumes that all individuals have the same number of contacts. The mass-action model could potentially find wide utility in populations that are heterogeneous with respect to contact rates, infectiousness or susceptibility, specifically for modelling highly contagious diseases (such that brief contacts lead to transmission) or slow-propagating infectious diseases (such as many STDs); in either case, epidemic propagation is slow relative to the turnover in contacts.

We check our mathematical results using simulations that model continuous-time stochastic processes (both social and epidemiological) and take into account the finite size and heterogeneity of the population (§3). We wish to highlight our specific simulation techniques as an interesting alternative to the commonly used chain-binomial simulation (Daley *et al.* 2001).

The NE model offers a flexible starting point for analysing epidemiological processes in dynamic networks. Although we have started by analysing a simple category of dynamic random network, it should be fairly straightforward to extend the model to populations with simple spatial heterogeneity or assortative mixing by type (Newman 2003). Indeed, similar methods for static networks have recently been extended to directed networks, networks with assortativity, and affiliation structure (Newman 2002b; Meyers & Newman 2003; Meyers *et al.* 2006). It is probable that these, and other extensions, can be made to the NE model.

The authors thank Richard Rothenberg for providing the Atlanta dataset and useful comments. L.A.M. acknowledges grant support from the James S. McDonnell Foundation.

REFERENCES

- Adimora, A. A., Schoenbach, V. J. & Doherty, I. A. 2006 HIV and African Americans in the southern United States: sexual networks and social context. *Sex. Transm. Dis.* **33**, S39–S45. (doi:10.1097/01.olq.0000228298.07826.68)
- Altmann, M. 1995 Susceptible–infected–removed epidemic models with dynamic partnerships. *J. Math. Biol.* **33**, 661–675. (doi:10.1007/BF00298647)
- Altmann, M. 1998 The deterministic limit of infectious disease models with dynamic partners. *Math. Biosci.* **150**, 153–175. (doi:10.1016/S0025-5564(98)00012-1)
- Anderson, R. & May, R. 1991 *Infectious diseases of humans: dynamics and control*. Oxford, UK: Oxford University Press.
- Andersson, H. 1998 Limit theorems for a random graph epidemic model. *Ann. Appl. Probab.* **8**, 1331–1349. (doi:10.1214/aoap/1028903384)
- Barthélemy, M., Barrat, A., Pastor-Satorras, R. & Vespignani, A. 2005 Dynamical patterns of epidemic outbreaks in complex heterogeneous networks. *J. Theor. Biol.* **235**, 275–288. (doi:10.1016/j.jtbi.2005.01.011)
- Bauch, C. 2002 A versatile ODE approximation to a network model for the spread of sexually transmitted diseases. *J. Math. Biol.* **45**, 375–395. (doi:10.1007/s002850200153)
- Boguna, M., Pastor-Satorras, R. & Vespignani, A. 2003 Epidemic spreading in complex networks with degree correlations. In *Statistical mechanics of complex networks*, vol. 625 (eds R. Pastor-Satorras, M. Rubi & A. Diaz-Guilera) *Lecture Notes in Physics*, pp. 127–147. Berlin, Germany: Springer.
- Callaway, D. S., Newman, M. E. J., Strogatz, S. H. & Watts, D. J. 2000 Network robustness and fragility: percolation on random graphs. *Phys. Rev. Lett.* **85**, 5468–5471. (doi:10.1103/PhysRevLett.85.5468)
- Chick, S. E., Adams, A. L. & Koopman, J. S. 2000 Analysis and simulation of a stochastic, discrete-individual model of STD transmission with partnership concurrency. *Math. Biosci.* **166**, 45–68. (doi:10.1016/S0025-5564(00)00028-6)
- Daley, D., Gani, J. & Gani, J. 2001 *Epidemic modelling: an introduction*. Cambridge, UK: Cambridge University Press.
- Diekmann, O. & Heesterbeek, J. 2000 *Mathematical epidemiology of infectious diseases. Model building, analysis and interpretation*. Chichester, UK: Wiley.
- Dietz, K. 1988 On the transmission dynamics of HIV. *Math. Biosci.* **90**, 397–414. (doi:10.1016/0025-5564(88)90077-6)
- Doherty, I. A., Shiboski, S., Ellen, J. M., Adimora, A. A. & Padian, N. S. 2006 Sexual bridging socially and over time: a simulation model exploring the relative effects of mixing and concurrency on viral sexually transmitted infection transmission. *Sex. Transm. Dis.* **33**, 368–373. (doi:10.1097/01.olq.0000194586.66409.7a)
- Eames, K. T. D. & Keeling, M. J. 2002 Modeling dynamic and network heterogeneities in the spread of sexually transmitted diseases. *Proc. Natl Acad. Sci. USA* **99**, 13 330–13 335. (doi:10.1073/pnas.202244299)
- Eames, K. T. D. & Keeling, M. J. 2004 Monogamous networks and the spread of sexually transmitted diseases. *Math. Biosci.* **189**, 115–130. (doi:10.1016/j.mbs.2004.02.003)
- Evans, T. S. & Plato, A. D. K. 2007 Exact solution for the time evolution of network rewiring models. *Phys. Rev. E* **75**, 056101. (doi:10.1103/PhysRevE.75.056101)
- Ferguson, N. M. & Garnett, G. P. 2000 More realistic models of sexually transmitted disease transmission dynamics: sexual partnership networks, pair models, and moment closure. *Sex. Transm. Dis.* **27**, 600–609. (doi:10.1097/00007435-200011000-00008)
- Finkenstadt, B. & Grenfell, B. 1998 Empirical determinants of measles metapopulation dynamics in England and Wales. *Proc. R. Soc. B* **265**, 211–220. (doi:10.1098/rspb.1998.0284)
- Ford, K., Sohn, W. & Lepkowski, J. 2002 American adolescents: sexual mixing patterns, bridge partners, and concurrency. *Sex. Transm. Dis.* **29**, 13–19. (doi:10.1097/00007435-200201000-00003)
- Garnett, G. P., Aral, S. O., Hoyle, D. V., Cates Jr, W. & Anderson, R. M. 1997 The natural history of syphilis: implications for the transmission dynamics and control of infection. *Sex. Transm. Dis.* **24**, 185–200.
- Grassly, N. C., Fraser, C. & Garnett, G. P. 2005 Host immunity and synchronized epidemics of syphilis across the United States. *Nature* **433**, 417–421. (doi:10.1038/nature03072)

- Grenfell, B. T., Bjørnstad, O. N. & Kappey, J. 2001 Travelling waves and spatial hierarchies in measles epidemics. *Nature* **414**, 716–723. (doi:10.1038/414716a)
- Gupta, S., Anderson, R. & May, R. 1989 Networks of sexual contacts: implications for the pattern of spread of HIV. *AIDS* **3**, 807–817. (doi:10.1097/00002030-198912000-00005)
- Heckathorn, D. 2002 Respondent-driven sampling ii: deriving valid population estimates from chain-referral samples of hidden populations. *Soc. Probl.* **49**, 11–34. (doi:10.1525/sp.2002.49.1.11)
- Holme, P. & Zhao, J. 2007 Exploring the assortativity-clustering space of a network's degree sequence. *Phys. Rev. E* **75**, 046111. (doi:10.1103/PhysRevE.75.046111)
- Jones, S. 2005 Analysis of a syphilis outbreak through the lens of percolation theory. Master's thesis, University of Texas-Austin.
- Liljeros, F., Edling, C. R., Amaral, L. A., Stanley, H. E. & Åberg, Y. 2001 The web of human sexual contacts. *Nature* **411**, 907–908. (doi:10.1038/35082140)
- Lloyd, A. L. & May, R. M. 1996 Spatial heterogeneity in epidemic models. *J. Theoret. Biol.* **179**, 1–11. (doi:10.1006/jtbi.1996.0042)
- Manhart, L. E., Aral, S. O., Holmes, K. K. & Foxman, B. 2002 Sex partner concurrency: measurement, prevalence, and correlates among urban 18–39-year-olds. *Sex. Transm. Dis.* **29**, 133–143. (doi:10.1097/00007435-200203000-00003)
- Maslov, S. & Sneppen, K. 2002 Specificity and stability in topology of protein networks. *Science* **296**, 910–913. (doi:10.1126/science.1065103)
- Meyers, L. A. & Newman, M. E. J. 2003 Applying network theory to epidemics: control measures for mycoplasma pneumoniae outbreaks. *Emerg. Infect. Dis.* **9**, 204.
- Meyers, L. A., Pourbohloul, B., Newman, M. E. J., Skowronski, D. M. & Brunham, R. C. 2005 Network theory and SARS: predicting outbreak diversity. *J. Theor. Biol.* **232**, 71–81. (doi:10.1016/j.jtbi.2004.07.026)
- Meyers, L. A., Newman, M. E. J. & Pourbohloul, B. 2006 Predicting epidemics on directed contact networks. *J. Theor. Biol.* **240**, 400–418. (doi:10.1016/j.jtbi.2005.10.004)
- Molloy, M. & Reed, B. 1995 A critical point for random graphs with a given degree sequence. *Random Struct. Algorit.* **6**, 161.
- Molloy, M. & Reed, B. 1998 The size of the giant component of a random graph with a given degree sequence. *Combin. Probab. Comput.* **7**, 295–305. (doi:10.1017/S0963548398003526)
- Newman, M. E. J. 2002a The spread of epidemic disease on networks. *Phys. Rev. E* **66**, 016128. (doi:10.1103/PhysRevE.66.016128)
- Newman, M. E. J. 2002b Assortative mixing in networks. *Phys. Rev. Lett.* **89**, 208701. (doi:10.1103/PhysRevLett.89.208701)
- Newman, M. E. J. 2003 Mixing patterns in networks. *Phys. Rev. E* **67**, 026126. (doi:10.1103/PhysRevE.67.026126)
- Newman, M. E. J., Watts, D. J. & Strogatz, S. H. 2002 Random graph models of social networks. *Proc. Natl Acad. Sci. USA* **99**, 2566–2572. (doi:10.1073/pnas.012582999)
- Pastor-Satorras, R. & Vespignani, A. 2001a Epidemic spreading in scale-free networks. *Phys. Rev. Lett.* **86**, 3200–3203. (doi:10.1103/PhysRevLett.86.3200)
- Pastor-Satorras, R. & Vespignani, A. 2001b Epidemic dynamics and endemic states in complex networks. *Phys. Rev. E* **63**, 066117. (doi:10.1103/PhysRevE.63.066117)
- Ross, R. 1910 *The prevention of malaria*. London, UK: John Murray.
- Rothenberg, R., Sterk, C., Toomey, K., Potterat, J., Johnson, D., Schrader, M. & Hatch, S. 1998 Using social network and ethnographic tools to evaluate syphilis transmission. *Sex. Transm. Dis.* **25**, 154–160. (doi:10.1097/00007435-199803000-00009)
- Salganik, M. J. & Heckathorn, D. D. 2004 Sampling and estimation in hidden populations using respondent-driven sampling. *Sociol. Methodol.* **34**, 193–240. (doi:10.1111/j.0081-1750.2004.00152.x)
- Strogatz, S. H. 2001 Exploring complex networks. *Nature* **410**, 268–276. (doi:10.1038/35065725)
- van den Driessche, P. & Watmough, J. 2002 Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Math. Biosci.* **180**, 29–48. (doi:10.1016/S0025-5564(02)00108-6)
- Volz, E. 2007 SIR dynamics in random networks with heterogeneous connectivity. *J. math. Biol.* (doi:10.1007/s00285-007-0116-4)
- Volz, E. & Heckathorn, D. In press. Probability based estimation theory for respondent driven sampling. *J. Official Statistics*.
- Watts, C. H. & May, R. M. 1992 The influence of concurrent partnerships on the dynamics of HIV/AIDS. *Math. Biosci.* **108**, 89–104. (doi:10.1016/0025-5564(92)90006-I)
- Watts, D. J. & Strogatz, S. H. 1998 Collective dynamics of small-world networks. *Nature* **393**, 409–410. (doi:10.1038/30918)
- Watts, D. J., Muhamad, R., Medina, D. C. & Dodds, P. S. 2005 Multiscale, resurgent epidemics in a hierarchical metapopulation model. *Proc. Natl Acad. Sci. USA* **102**, 11157–11162. (doi:10.1073/pnas.0501226102)

NOTICE OF CORRECTION

Equations (2.17), (2.18) and [table 4](#) are now presented in the correct form.

28 May 2008